READ ENCLOSED LEAFLET BEFORE USE.

INDOFIL INDUSTRIES LIMITED

Mesotrione 2.27% + Atrazine 22.7% SC

Herbicide

Marketed by :
Indofil Industries Limited
Regd. Office: Kalpataru Square, 4th Floor, Kondivita Road, Off Andheri-Kurla Road, Andheri (East), Mumbai - 400059

विपणनकर्ता : इंडोफिल इण्डस्ट्रीज लिमिटेड
रजि. ऑफिस: कल्पतरु स्क्वेअर, ४थी मंजिल, कोन्डीविटा रोड, ऑफ अंधेरी–कुर्ला रोड, अंधेरी (पूर्व), मुंबई – ४०००५९

Manufactured by:
Parijat Industries (India) Pvt. Ltd.
Office: M-77, Ist and IInd Floor, M-Block Market, Greater Kailash- II. Factory: Unit – 1 Village Mandhaur, Naraingarh Road, Ambala, Haryana.
Unit – 2 Village Khera Gani, near Village Fatehgarh, Raipur Rani Road, Distt. Ambala (Haryana).

For any complaint contact Customer Care Cell at Indofil Industries Limited, Kalpataru Square, 4th Floor, Kondivita Road, Off Andheri-Kurla Road, Andheri (East), Mumbai - 400 059. Toll Free No.: 1800-120-003-004, email: customercare@indofil.com

Not to be used post Harvest application
हार्वेस्ट आवेदन पर उपयोग के लिए नहीं
कापणी अनुप्रयोग पोस्ट वापरले जाऊ नाही
હાર્વેસ્ટ અરજી પોસ્ટ વાપરી શકાય નથી
ਵਾਢੀ ਐਪਲੀਕੇਸ਼ਨ ਤੋਂ ਪੋਸਟ ਵਰਤਿਆ ਜਾ ਨਾ ਕਰਨ
ஹார்வெஸ்ட் ஆப்ளிகேஷன் பொஸ்ட் உபயோகிக்கப் படாது
హార్వెస్ట్ అప్లికేషన్ పోస్ట్ ఉపయోగించరాదు
ಹಾರ್ವೆಸ್ಟ್ ಅರ್ಜಿಯನ್ನು ಉಪಯೋಗಿಸಲು ಅಲ್ಲ
ഹാർവെസ്റ്റ് ആപ്ലിക്കേഷ പോസ്റ്റ് ഉപയോഗിക്കാ പാടില്ല
খালি কৰ্ত্তনৰ্স পুনৰব্যৱহাৰ বিচাঁ জনক

Warranty: Since the use of the product is beyond our control we cannot assume any responsibility other than for the uniform quality of the product.

For Agriculture use only. केवल कृषि उपयोग के लिए।



Mesotrione 2.27% + Atrazine 22.7% SC

Herbicide

READ ENCLOSED LEAFLET BEFORE USE.

INDOFIL INDUSTRIES LIMITED

Mesotrione 2.27% + Atrazine 22.7% SC

Herbicide

Marketed by :
Indofil Industries Limited
Regd. Office: Kalpataru Square, 4th Floor, Kondivita Road, Off Andheri-Kurla Road, Andheri (East), Mumbai - 400059

विपणनकर्ता : इंडोफिल इण्डस्ट्रीज लिमिटेड
रजि. ऑफिस: कल्पतरु स्क्वेअर, ४थी मंजिल, कोन्डीविटा रोड, ऑफ अंधेरी–कुर्ला रोड, अंधेरी (पूर्व), मुंबई – ४०००५९

Manufactured by:
Parijat Industries (India) Pvt. Ltd.
Office: M-77, Ist and IInd Floor, M-Block Market, Greater Kailash- II. Factory: Unit – 1 Village Mandhaur, Naraingarh Road, Ambala, Haryana.
Unit – 2 Village Khera Gani, near Village Fatehgarh, Raipur Rani Road, Distt. Ambala (Haryana).

For any complaint contact Customer Care Cell at Indofil Industries Limited, Kalpataru Square, 4th Floor, Kondivita Road, Off Andheri-Kurla Road, Andheri (East), Mumbai - 400 059. Toll Free No.: 1800-120-003-004, email: customercare@indofil.com

Not to be used post Harvest application
हार्वेस्ट आवेदन पर उपयोग के लिए नहीं
कापणी अनुप्रयोग पोस्ट वापरले जाऊ नाही
હાર્વેસ્ટ અરજી પોસ્ટ વાપરી શકાય નથી
ਵਾਢੀ ਐਪਲੀਕੇਸ਼ਨ ਤੋਂ ਪੋਸਟ ਵਰਤਿਆ ਜਾ ਨਾ ਕਰਨ
ஹார்வெஸ்ட் ஆப்ளிகேஷ பொஸ்ட் உபயோகிக்கப் படாது
హార్వెస్ట్ అప్లికేషన్ పోస్ట్ ఉపయోగించరాదు
ಹಾರ್ವೆಸ್ಟ್ ಅರ್ಜಿಯನ್ನು ಉಪಯೋಗಿಸಲು ಅಲ್ಲ
ഹാർവെസ്റ്റ് ആപ്ലിക്കേഷ പോസ്റ്റ് ഉപയോഗിക്കാ പാടില്ല
খালি কৰ্ত্তনৰ্স পুনৰব্যৱহাৰ বিচাঁ জনক

Warranty: Since the use of the product is beyond our control we cannot assume any responsibility other than for the uniform quality of the product.

For Agriculture use only. केवल कृषि उपयोग के लिए।



Mesotrione 2.27% + Atrazine 22.7% SC

Herbicide

GADGET

Mesotrione 2.27%+ Atrazine 22.7% SC (Selective Herbicide)

Introduction: Mesotrione 2.27%w/w + Atrazine 22.7%w/w SC is a selective post emergence herbicide recommended for the control of Grasses, Broad leaf weeds and Sedges in Maize and Sugarcane.

Crops and Pests group,Dosage:

Crop (s)	Common name of Weeds	Dosage/HA A.I (g)	Dosage/HA Formulation (ml)	Dilution in Water (L)	Waiting period (Days)	Re-entry after each application (In Hours)
Maize	Trianthema spp. Cyperus spp. Digitaria sanguinalis Echinochloa spp. Dactyloctenium aegyptium	875	3500	500	42	–
Sugar-cane	Trianthema spp. Amaranthus viridis Echinochloa colona Digitaria sanguinalis Cyperus rotundus Dactyloctenium aegyptium	875	3500	500	190	–

Manner and time of application: Apply as post emergence single spray application at 3-4 leaf stage of actively growing weeds. **Method of Application: Instructions for mixing:** Measure out the required quantity of the product and mix it well with a small quantity of water. Add the remaining quantity of water as specified with thorough agitation. Adjust the quantity of product in proportion to the water required to cover a hectare and apply by a suitable sprayer. **Symptoms of poisoning:** Irritation to eyes, skin and respiratory tract may occur, product is skin sensitizer so it may cause skin allergy and other non-specific symptoms may be expected as labored respiration, nausea, vomiting, diarrhoea, headache and confusion. **First Aid:** If poisoning or any indisposition is suspected. Stop work immediately and call a physician. Show label to the physician. Remove the affected person from danger zone to a well-ventilated room or to fresh air. If Ingested: Never give anything by mouth to an unconscious patient and never induce vomiting. Consult the doctor immediately. If Inhaled: Take the patient to fresh air and take care of body temperature. If patient is not breathing or breathing with difficulty, give artificial respiration, preferably with mouth-to-mouth breathing. Consult a physician. In case of skin contact: Affected parts of body e.g. finger, nails, etc should be washed with soap and water. All the contaminated clothes should be segregated and thoroughly washed with soap and water separately before use. If skin irritation persists, consult a doctor. In case of eye contact: Rinse gently with plenty of water for 15 minutes. If pain, irritation, redness, or photophobia persists, consult eye specialist. **Antidote:** No specific antidote is known; apply symptomatic therapy. **Phytotoxicity:** Not phytotoxic to crop at recommended dosage. **Storage:** 1. The packages containing the herbicide should be tightly closed, stored in original containers in separate rooms or premises away from the rooms or premises used for storing other articles particularly articles of food and animal feed or shall be kept in separate almirahs under lock and key, away from children, unauthorized persons and animals. 2. The rooms or premises meant for storing the herbicide shall be well built, dry, well-lit and ventilated and of sufficient dimension to avoid contamination with vapor. Keep away from direct sunlight. 3. Do not allow pilferage of stored product. **Disposal of used container:** 1. It shall be the duty of the manufacturers, formulators of insecticide and operators to dispose packages or surplus materials and washings in a safe manner so as to prevent environmental or water pollution. 2. The used packages shall not be left outside to prevent misuse. 3. Packages shall not be broken and buried away from habitations. **Dosage/Area (Per Hectare):** As mentioned on table. **Weather Condition (in general):** Do not spray during the windy weather conditions or during the rainy days, if the foliage is wet or if rain is imminent. Spray strictly in the direction of w to prevent prolonged or repeated exposure to the spray mist. Do not under dose or overdose overlapping applications or by exceeding the recommended rate. **Soil & Water (Wherever applicable):** This product should be applied during optimum sufficient soil moisture conditions. Do not contaminate water used for irrigation or domestic purpose, avoid spray near water areas viz. ponds, ditches, lakes, drainage systems, etc. by disposal of product waste and do not allow direct over spray or spray drift of product on these areas to avoid long term adverse effect in the aquatic environment. **Application Techniques:** Spray the recommended dose of this herbicide as post emergence single spray application at 3-4 leaf stage of actively growing weeds by using sufficient quantity of water to ensure thorough coverage of the weed foliage with the help of knapsack or any other suitable sprayer fitted with flat fan or flood jet deflector nozzle. **Application Timings:** Apply as post emergence single spray application at 3-4 leaf stage of actively growing weeds. **Crop Stage For Application:** Spray the recommended dose of this herbicide as post emergence single spray application at 3-4 leaf stage of actively growing weeds by using sufficient quantity of water to ensure thorough coverage of the weed foliage. **Equipment type:** Spray with knapsack sprayer or any other suitable sprayer. **Nozzle type:** Flat fan or flood jet deflector nozzle. **Optimal Re-entry period:** Do not enter treated areas for 24 hours after spray application. **Pre-Harvest Interval (Days):** Mentioned in the table. **Frequency Including max Number of Sprays:** Frequency including maximum number of sprays: Apply as post emergence single spray application at 3-4 leaf stage of actively growing weeds. Maximum no. of applications is one. **Purpose:** for Agricultural Use. **Composition of Herbicides:** Mesotrione A.I. 2.27%w/w, Atrazine A.I. 22.7%w/w, Xanthan gum (Thickener) 0.14%w/w, Attapulgite clay (Thickener) 0.9%w/w, 1,2-benzisothiazol-3-one (as a 20% solution) 0.18%w/w, Polydimethyl siloxane (Antifoam agent) 0.43%w/w, Citric acid (Buffers) 0.91%w/w, Phosphoric acid solution (Buffers) 1.0 %w/w, Aqueous solution of D-glucopyranose oligomeric C9- C11 alkyl glycosides (Adjuvant) 18.18%w/w, Water (Solvent) Q.S.%w/w, Acrylic graft copolymer solution (Dispersing Agent) 1.82%w/w, Isotridecanol-8-ethoxylate (Dispersing Agent) 1.82%w/w, 1,2-propanediol (Antifreeze) 4.55%w/w, Tetrasodium diphosphate (Stabilizing agent) 0.03%w/w, **Total: 100%w/w. Precautions:** Do not use cooking utensils for preparing the spray solution. Use wooden stick for stirring the spray solution. Avoid contact with skin, eyes, mouth and clothing. Avoid inhalation of fog, vapors or spray mist. Wear protective clothing (full sleeve shirt and pants), chemical resistant hand gloves to avoid contact with the skin, mask, goggles, hat and boots, while handling the product. Do not eat, drink or smoke while applying the product. Wash hands with soap and plenty of water and change clothes after the work is over. Avoid contamination of air and water bodies with the herbicide. Also wash the contaminated clothes before re-use. Newly treated areas must not be grazed and all livestock is to be kept out until the treated area is dry. This product should not be applied (tank – mix) with any other chemical. Do not blow nozzle with mouth. Do not spray crops under stress. If any indisposition occurs, stop the work, follow first aid measures and call a physician immediately. Spilt product must not be re-used. Collect up spilt material thoroughly and bury it in a safe place. Rinse spilt area with plenty of water. It is dangerous to re-use empty containers. Rinse empty container three times before disposal and add the rinsed into spray tank before application. Do not re-use empty container to keep food items or for any other purpose. **Cautionary statement: 1. Not to be used for any purpose other than those subscribed on label and leaflets 2. Destroy the container after use as directed on leaflet. 3. Toxic to fish.**

ENGLISH

गॅजेट

मिजोट्राइन २.२७% + एट्राजीन २२.७% एस. सी. (चयनात्मक खरपतवारनाशक)

परिचय: मिजोट्राइन २.२७% भार/भार + एट्राजीन २२.७% भार/भार एस.सी.एक चुनिंदा खरपतवारनाशक है जिसका इस्तेमाल मक्का एवं गन्ना फसलों में घासकुल वाले, चौडी पत्तीवाले जाती के खरपतवारों एवं सेज के नियंत्रण हेतु खरपतवार उगने के बाद किया जाता है। चेतावनियाँ: १. इस खरपतवारनाशक का लेबल पर दिये हुए विवरण के अनुसार इस्तेमाल करने के अलावा किसी अन्य उद्देश्य हेतु इस्तेमाल नहीं करें। २. यह उत्पाद मछली और जलीय जीवों के लिए विषाक्त है, इसका उपयोग जलीय कृषि /मत्स्य पालन के आसपास ना करें। ३. उत्पाद त्वचा संवेदी है, लंबे समय तक या बार–बार त्वचा के संपर्क में आने से एलर्जी हो सकती है।

फसलें और कीट समूह, मात्रा:

फसल	खरपतवार का नाम	प्रति हेक्टेअर मात्रा स.तत्व (ग्रा.)	प्रति हेक्टेअर मात्रा संरचना (मि.लि.)	पानी की मात्रा (ली.)	अन्तिम छिडकाव तथा फसल काटने के बीच अन्तराल (दिन)	पुन: प्रवेश की अवधि प्रत्येक छिडकाव के बाद (घंटे)
मक्का	इटसीट (ट्राअमन्थीसीव), मोथा (साइप्रसएसीसी), बनशाप (डिजिटेरियासैंगुइनलिस), सॉवा (एचिनोक्लोआएसीयर), नकरा घास (डक्टीलोक्टेनियम एजिप्टेवम)	८७५	३५००	५००	४२	–
गन्ना	इटसीट (ट्राअमन्थीसीव), चौलइ (अमरांथसविरिडेस), सॉवा (एचिनोक्लोआकोलोना), बनशाप (डिजिटेरियासैंगुइनलिस), बज्ञानगलोथ(साइप्रसरोटंडस), नकराघास (डक्टीलोक्टेनियमएजिप्टेवम)	८७५	३५००	५००	१९०	–

रीति और अनुप्रयोग का समय: खरपतवारों के ३–४ पत्ती की सक्रिय रूप से उगने की अवस्था होने पर केवल एक बार। **उपयोग की दिशा और इस्तेमाल का तरीका:** इस्तेमाल के लिए निर्देश उत्पाद की आवश्यक मात्रा को नापें और इसे थोडी मात्रा में पानी के साथ अच्छी तरह मिलाएं। पूरी तरह से हिलाने के साथ निर्दिष्ट पानी की शेष मात्रा जोड़ें तथा किसी योग्य स्प्रेयर से पूरी फसल में छिड़काव करें और उपयोग करते समय ओवर लैपिंग द्वारा या अनुशंसित दर से कम /अधिक उपयोग के कारण मात्रा में छिड़काव न करें। **विष के लक्षण:** आंखों, त्वचा और श्वसन नलिका में जलन हो सकती है, उत्पाद त्वचा संवेदनशील है, इसलिए यह त्वचा एलर्जी का कारण हो सकता है और अन्य गैर–विशिष्ट लक्षणों को संभव माना जा सकता है जैसे श्वासोच्छवास, मळमळ, उलटया, अतिसार, डोकेदुखी आणि भ्रम. **प्राथमिक चिकित्सा:** विषबाधा किंवा कोणत्याही असस्वस्थतेचा संशय असल्यास तात्काळीक काम थांबवा आणि डॉक्टरांना कॉल करा. डॉक्टरांना लेबल दाखवा. बाधित व्यक्तीला धोक्याच्या क्षेत्रातून हवेशीर वातीनि किंवा ताज्या हवेत न्या. गिळणे: बेशुद्ध व्यक्तीला कधीही काही तोंडात घालू नका. उलट्या करू नका. डॉक्टरांचा सल्ला घ्या. श्वसन: बाधित व्यक्तीला ताजी हवा द्या. त्वचा: त्वचेवर पाणी आणि साबण वापरा. डोळे: डोळ्यात थेट फवारणी टाळा. डोळ्यात फवारणी झाल्यास ताबडतोब डोळ्यात स्वच्छ पाणी घाला. **विषारीपणा:** पिकांसाठी शिफारशीनुसार वापरल्यास पिकांसाठी विषारी नाही. **साठवण:** १. कीटकनाशक असलेले पॅकेजेस घट्ट बंद करून, मूळ कंटेनरमध्ये वेगळ्या खोल्यांमध्ये किंवा अन्नासह साठवून ठेवलेली पावडरसह किंवा इतर वस्तू विशेषतः अन्न आणि पशुखाद्य साठवण्यासाठी वापरल्या जाणाऱ्या खोल्या किंवा परिसरापासून दूर ठेवावयास किंवा कुलूपबंद स्वतंत्र कपाटांमध्ये ठेवाव्यात आणि मुले, अनधिकृत व्यक्ती आणि प्राणी यांच्यापासून दूर ठेवावे. २. तणनाशक साठवण्यासाठी असलेल्या खोल्या किंवा जागा चांगल्या बांधलेल्या, कोरड्या, चांगले प्रकाशित आणि हवेशीर आणि बाष्पाने दूषित होऊ नये म्हणून पुरेशा आकाराचे असावेत. थेट सूर्यप्रकाशापासून दूर ठेवा. ३. उत्पादन गोतवू देऊ नका. **वापरलेल्या कंटेनरची विल्हेवाट लावणे:** १. पर्यावरण किंवा जल प्रदूषण रोखण्यासाठी पॅकेजेस किंवा अतिरिक्त साहित्य आणि वॉशिंगची सुरक्षितपणे विल्हेवाट लावणे हे उत्पादक, कीटकनाशकांचे सूत्रकार आणि ऑपरेटर यांची सर्वस्व कर्तव्य असेल. २. वापरलेले पॅकेज त्यांचा पुन्हा वापर टाळण्यासाठी बाहेर ठेऊ नये. ३. पाकिटे तोडून वस्तीपासून दूर पुरले जावे. **मात्रा/क्षेत्र (प्रति हेक्टर):** कोष्टकात नमूद केल्याप्रमाणे. **हवामानाची स्थिती (सर्वसाधारणपणे):** वाऱ्याच्या वातावरणात किंवा दिवसा गरम असताना, पर्णसंभार ओला असल्यास किंवा पाऊस जवळ येत असल्यास फवारणी करू नका. फवारणी धुकेचा दीर्घकाळ किंवा वारंवार संपर्क टाळण्यासाठी वाऱ्याच्या दिशेने काटेकोरपणे फवारणी करा. कमी मात्रा किंवा अतिमात्रा एकमेकांवर फवारणी किंवा शिफारस केलेल्या दरापेक्षा जास्त फवारणी करू नका. **माती आणि पाणी (जिथे लागू असेल):** हे उत्पादन इष्टतम पुरेशा जमिनीतील आर्द्रतेच्या परिस्थितीत लागू केले पाहिजे. सिंचन किंवा घरगुती कारणासाठी वापरलेले पाणी दूषित करू नका आणि पाण्याचे क्षेत्र जसे. तलाव, खड्डे, तलाव, सांडपाणी व्यवस्था इ. उत्पादनाच्या कचऱ्याची विल्हेवाट लावा आणि जलीय वातावरणावर दीर्घकालीन प्रतिकूल परिणाम टाळण्यासाठी या भागात थेट ओव्हर स्प्रे किंवा फवारणी उत्पादनास वाहून जाऊ देऊ नका. **वापरण्याचे तंत्र:** या तणनाशकाची शिफारस केलेल्या मात्रेची फवारणी एकच फवारणी म्हणून ३–४ पानांच्या अवस्थेवर सक्रियपणे वाढणाऱ्या तणांच्या पानांवर पाण्याच्या टप्प्यावर करा, जेणेकरून नॅपसॅक किंवा फ्लॅट फॅन किंवा फ्लड जेट डिफ्लेक्टर नोजलसह बसवलेल्या इतर कोणत्याही योग्य स्प्रेअरने फवारणी यंत्राच्या सहाय्याने तणांच्या पानांचे संपूर्ण कव्हरेज सुनिश्चित करण्यासाठी पुरेशा प्रमाणात पाण्याचा वापर करा. **वापराच्या वेळा:** सक्रियपणे वाढणाऱ्या तणांच्या ३–४ पानांच्या अवस्थेत सक्रियपणे वाढणारी तणांच्या पानांच्या टप्प्यावर करा जेणेकरून नॅपसॅक किंवा फ्लॅट फॅन किंवा फ्लड जेट डिफ्लेक्टर नोजलसह बसवलेल्या इतर कोणत्याही योग्य स्प्रेअरने फवारणी करा. **नोजल प्रकार:** फ्लॅट फॅन किंवा फ्लड जेट डिफ्लेक्टर नोजल. **इष्टतम पुनर्प्रवेश कालावधी:** फवारणी केल्यानंतर 24 तास उपचार केलेल्या भागात प्रवेश करू नका. **काढणीपूर्व अंतर (दिवस):** तक्त्यामध्ये नमूद केले आहे. **फवारणीच्या कमाल संख्येसह वारंवारता:** जास्तीत जास्त फवारण्यासह वारंवारता: सामान्य असलेली वाढणाऱ्या तणांच्या ३–४ पानांच्या टप्प्यावर एकच फवारणी म्हणून उगवलेल्यांवर करा. फवारणीची कमाल संख्या एक आहे. **उद्देश:** शेतीसाठी वापरलेले पाणी दूषित करू नका आणि पाण्याचे क्षेत्र जसे. तलाव, खड्डे, तलाव, सांडपाणी व्यवस्था इ. उत्पादनाच्या कचऱ्याची विल्हेवाट लावून पर्यावरण किंवा पाण्याचे प्रदूषण टाळा आणि उत्पादनाच्या थेट फवारणी किंवा फवारणी टाळा जेणेकरून जलीय वातावरणात दीर्घकालीन प्रतिकूल परिणाम टाळता येईल. **अर्जाचे तंत्र:** या तणनाशकाची शिफारस केलेल्या मात्रेची फवारणी एकच फवारणी म्हणून ३–४ पानांच्या अवस्थेत सक्रियपणे वाढणाऱ्या तणांच्या पानांवर पाण्याच्या टप्प्यावर करा. **पीक टप्पा अर्जासाठी:** या तणनाशकाची शिफारस केलेल्या मात्रेची फवारणी एकच फवारणी म्हणून ३–४ पानांच्या अवस्थेत सक्रियपणे वाढणाऱ्या तणांच्या पानांवर पाण्याच्या टप्प्यावर करा. **उपकरणाचा प्रकार:** नॅपसॅक स्प्रेअर किंवा इतर कोणत्याही योग्य स्प्रेअरने फवारणी करा. **नोजल प्रकार:** फ्लॅट फॅन किंवा फ्लड जेट डिफ्लेक्टर नोजल. **इष्टतम पुनर्प्रवेश कालावधी:** फवारणी केल्यानंतर 24 तास उपचार केलेल्या भागात प्रवेश करू नका. **काढणीपूर्व अंतर (दिवस):** तक्त्यामध्ये नमूद केले आहे. **फवारणीच्या कमाल संख्येसह वारंवारता:** जास्तीत जास्त फवारण्यासह वारंवारता: सामान्य असलेली वाढणाऱ्या तणांच्या ३–४ पानांच्या टप्प्यावर एकच फवारणी म्हणून उगवलेल्यांवर करा. फवारणीची कमाल संख्या एक आहे. **उद्देश:** शेतीसाठी वापरलेले पाणी दूषित करू नका आणि पाण्याचे क्षेत्र जसे. तलाव, खड्डे, तलाव, सांडपाणी व्यवस्था इ. उत्पादनाच्या कचऱ्याची विल्हेवाट लावून पर्यावरण किंवा पाण्याचे प्रदूषण टाळा आणि उत्पादनाच्या थेट फवारणी किंवा फवारणी टाळा जेणेकरून जलीय वातावरणात दीर्घकालीन प्रतिकूल परिणाम टाळता येईल. **अर्जाचे तंत्र:** या तणनाशकाची शिफारस केलेल्या मात्रेची फवारणी एकच फवारणी म्हणून ३–४ पानांच्या अवस्थेत सक्रियपणे वाढणाऱ्या तणांच्या पानांवर पाण्याच्या टप्प्यावर करा. **पीक टप्पा अर्जासाठी:** या तणनाशकाची शिफारस केलेल्या मात्रेची फवारणी एकच फवारणी म्हणून ३–४ पानांच्या अवस्थेत सक्रियपणे वाढणाऱ्या तणांच्या पानांवर पाण्याच्या टप्प्यावर करा. **उपकरणाचा प्रकार:** नॅपसॅक स्प्रेअर किंवा इतर कोणत्याही योग्य स्प्रेअरने फवारणी करा. **नोजल प्रकार:** फ्लॅट फॅन किंवा फ्लड जेट डिफ्लेक्टर नोजल. **पुनःप्रवेशाचा इष्टतम समयावधी:** हेक्टर छंटकाव पछी 24 हवाले सुधी छंटकाव करेला विस्तारमां प्रवेश करवो जोईये. **कापणी पहेलांनो समयावधी (दिवसो):** कोष्टकमां दर्शावेल मुजब. **वधुमां वधु वार छंटकाव सहित छंटकावनी आवृत्ति:** वधुमां वधु वार छंटकाव सहित छंटकावनी आवृत्ति: सक्रिय रीते वधतां नदीन तणो पर एकवार ३–४ पानी स्थितिमां सांच त्यारे એકવાર છંટકાવ કરો. **વધુમાં વધુ વાર છંટકાવ સહિત છંટકાવની આવૃત્તિ:** વધુમાં વધુ વાર છંટકાવ સહિત છંટકાવની આવૃત્તિ: સક્રિય રીતે વધતું નદીન તણ પર ૩–૪ પાનની સ્થિતિમાં સાંચ ત્યારે