HOKORI®
DINOTEFURAN 5% + ETHION 50% EC
(Contact and systemic Insecticide)
For Agricultural use only

Dinotefuran 5% + Ethion 50% (55 % w/w) EC is an emulsifiable concentration which is contact and systemic in nature. The insecticide exhibits translaminar activity and is recommended for the control of Brown Plant hopper, White backed Plant Hopper, Green leaf hopper and Leaf folder of rice crop.

Caution: 1. Product is toxic to fish and honey bees, it should not be used in the area where pisciculture/aquaculture are practiced and during foraging period of honey bees. 2. Destroy the container after use as directed in the leaflet.

Chemical Composition:

Ingredient	Content (% by w/w)
Dinotefuran Technical (a.i.)	5.00
Ethion Technical (a.i.)	50.00
Aromatic Alcohol (Co-Solvent)	6.00
Cyclic Carboxylic Esters (Solvent)	Q.S
Epoxide of Polyunsaturated Vegetable Alcohol (Stabilizer)	3.00
Alkali Metal Salt of Alkyl Aryl Sulphonic Acid (Emulsifier)	2.20
Blend of Polyoxy Ethylene Alkyl Aryl Alcoholic Ether (IGSURF -5050)	5.60
Total	**100.00**

Application & recommendation for use: When used as per the recommendations given below, Dinotefuran 5% + Ethion 50% (55 % w/w) EC provides effective management of Brown Plant hopper, White backed plant hopper, Green leaf hopper and Leaf folder of rice crop.

Crop	Common Name of the Pest	Dosage per Ha			Waiting period from last application to harvest
		a.i. (g)	Formulation (ml)	Dilution in water (L)	
Rice	Brown plant hopper, White backed Plant hopper, Green leaf hopper and Leaf folder	50+500	1000	500	33 days

Time of application and crop stage: Apply as soon as the pest population reaches at Economic Threshold Level (ETL).

Weather conditions: Should be applied preferably in the morning and evening. Avoid applications under high wind and heavy rains forecasted.

Soil: This is foliar application insecticide. So, hardly have any relation to soil.

Water: Clean water to be used for spray.

Application timings: Should be applied preferably in the morning and evening.

Application technique, equipment type, nozzle type: The product can be sprayed with all kind of high and low volume sprayers. It can be applied by manual and tractor mounted sprayer equipped with hollow cone nozzle using sufficient quantity of water to ensure thorough coverage.

Dosage: 1000 ml of formulation in 500 liters of water per hectare.

Instruction for mixing: Shake the bottle well before use. Measure out requisite quantity of the premix insecticide; add to it a small quantity of water. Stir thoroughly using a wooden stick. Add the remaining quantity of water as specified with thorough agitation to form final stock solution.

Re-entry period: 48 hours after spray.

Frequency including maximum number of sprays: Maximum 2-4 sprays depending upon target insect pest pressure.

Stages during which insecticide should not be applied on any crop: No restriction for crop stage. Follow the PHI recommendation for each crop. To be used as per the recommendation to control BPH, WBHP, GLH and Leaf folder of rice crop.

Usage of Pesticides in confined area: The product should be used as per the recommendation table.

Phytotoxicity: The product is not phytotoxic when used as per field recommendation.

Storage Conditions: 1. The package containing the insecticide shall be stored in separate rooms or premises away from the rooms or premises used for storing other articles particularly articles of food or shall be kept in separate almirahs under lock and key depending upon the quantity and nature of the insecticide. 2. The rooms or premises meant for storing the insecticide should be well built dry, well ventilated and of sufficient dimensions.

Safety Precautions: 1. Keep away from food, empty foodstuff containers and animal feed. 2. Avoid contact with mouth, skin and eyes. 3. Avoid inhalation of the spray mist. 4. Wash thoroughly the contaminated clothes and body parts after spraying. 5. Do not smoke, drink, eat and chew anything while spraying. 6. Use protective clothing which includes long sleeved shirt and long pants, chemical resistant gloves, goggles, shoes and socks while spraying.

Symptoms of Poisoning: Hypoactivity, staggered gait, Prostration, Miosis, Lacrimation, Excessive Salivation, Tachypnea, Dyspnoea, Convulsion & Tremors & soft stool.

First Aid - In case of inhalation: Immediately move victim to fresh air. If breathing is irregular or stopped, administer artificial respiration. Keep patient warm and at rest. Call doctor immediately. **In case of ingestion:** If swallowed, seek medical advice immediately. Do not induce vomiting. Never give anything by mouth to an unconscious person. **In case of skin contact:** Take off all contaminated clothing immediately and thoroughly wash the affected parts of the body with plenty of water. Wash contaminated clothes before re-use. **In case of eye contact:** Rinse immediately with plenty of water, also under the eyelids for atleast 15 minutes. If pain or irritation persists, consult eye specialist.

Advice to physician (Antidote): No specific antidote is known. Treat symptomatically.

Warning: Dangerous to reuse empty containers.

Disposal of Used Containers: 1. It shall be the duty of manufacturer, formulators of insecticides and operators to dispose off the washings and surplus materials in a safe manner so as to prevent environmental or water pollution. 2. The used packages shall not be left outside to prevent their reuse. 3. The packages shall be broken and buried away from habitation.

Manufactured by: ***PI Industries Ltd***
Regd. Office: Udaisagar Road, Udaipur-313001 (Rajasthan)
Unit 2: Plot No. 237, G.I.D.C. Industrial Estate, NH-8, Dist.: Bharuch, Panoli-394116, Gujarat
Unit 3: Plot No. 3133 to 3139 and others, G.I.D.C., Dist.: Bharuch, Panoli-394116, Gujarat
AN ISO 9001, 14001, 45001 & 27001

Marketed by: **Indofil Industries Limited.**
Regd. Office: Kalpataru Square, 4th Floor, Kondivita Road,
Off Andheri-Kurla Road, Andheri (East), Mumbai - 400 059

होकोरी
डिनोटेफुरान ५% + इथियॉन ५०% ई.सी.
(संपर्क और अंतःप्रवाही कीटनाशक)
केवल कृषि उपयोग हेतु

હોકોરી
ડિનોટેફુરાન 5% + ઈથિઓન 50% ઈ.સી.
(સંપર્ક અને આંતરપ્રવાહી જંતુનાશક)

होकोरी
डायनोटेफ्युरॉन 5% + इथियॉन 50% ई.सी.
(संपर्क आणि आंतरप्रवाही कीटकनाशक)
फक्त कृषी वापरासाठी

హోకోరి

ஹொக்கோரி
டைனோடெஃப்யூரான் 5% + எதியான் 50% EC

INSTRUCTION FOR USE

HOKORI®
DINOTEFURAN 5% + ETHION 50% EC
(Contact and systemic Insecticide)
For Agricultural use only

CUT HERE FOR DIRECTIONS FOR USE

ಹೊಕೊರಿ
ಡಿನೋಟೆಫುರನ್ 5% + ಎಥಿಯಾನ್ 50% ಇಸಿ
(ಸಂಪರ್ಕ ಮತ್ತು ಸಿಸ್ಟಮಿಕ್ ಕೀಟನಾಶಕ)
ಕೃಷಿ ಬಳಕೆಗೆ ಮಾತ್ರ

ਹੋਕੋਰੀ
ਡਾਇਨੋਟੇਫੂਰਾਨ 5% + ਇਥਿਓਨ 50% ਈ.ਸੀ
(ਸੰਪਰਕ ਅਤੇ ਅੰਤਰਪ੍ਰਵਾਹੀ ਕੀਟਨਾਸ਼ਕ)

হোকোরি
ডায়নোটেফিউরান ৫% + ইথিয়ন ৫০% ইসি

ହୋକୋରି

হোকৰী
ডিনোটেফুৰান 5% + ইথিয়ন 50% ইচি

BOOKLET PLACE HERE