# ZINEB 75% WP

# INDOFIL

## Z-78®

### CONTACT FUNGICIDE

जायनेब ७५ प्रतिशत समिश्रणीय पावडर

## इंडोफिल झेड-७८

(स्पर्शीय फफूंदीनाशक)

**Manufactured & Marketed by:**
**Indofil Industries Limited,**
**Regd. Office :** Kalpataru Square, 4th Floor, Kondivita Road, Off Andheri-Kurla Road, Andheri (East), Mumbai - 400 059

**उत्पादक एवं विक्रेता :**
**इंडोफिल इण्डस्ट्रीज लिमिटेड,**
**रजि.ऑफिस :** कल्पतरू स्क्वेअर, ४थी मंजिल, कोन्डीविटा रोड, ऑफ अन्धेरी–कुर्ला रोड, अन्धेरी (पूर्व), मुंबई–४०० ०५९

**Factory : Unit C -** D2/CH12, GIDC Estate Dahej, Tal. Vagra, Dist. Bharuch -392130, Gujarat. India
**Unit G -** Plot No. 2201, 2202, 2336, 2337, G.I.D.C. Panoli, Tal. - Ankleshwar, Dist.- Bharuch, Gujarat - 394116

**फैक्टरी : युनिट सी –** डी२/सीएच१२, जीआयडीसी इस्टेट दहेज, ता. वाग्रा, जि. भरुच–३९२१३०, गुजरात. भारत
**युनिट जी –** प्लॉट नं. २२०१, २२०२, २३३६, २३३७, जी.आय.डी.सी. पानोली, ता. – अंकलेश्वर, जि.– भरूच, गुजरात – ३९४११६

®-Registered Trademark of INDOFIL INDUSTRIES LIMITED, Mumbai, India.

For any complaint contact Customer Care Cell at Indofil Industries Limited, Kalpataru Square, 4th Floor, Kondivita Road, Off Andheri-Kurla Road, Andheri (East), Mumbai - 400 059.
Toll Free No. : 1800-120-003-004,
email : customercare@indofil.com

**For Identification of Factory please see first Alphabet of Batch No.**
**फैक्टरी की पहचान के लिए बैच नंबर का पहला अक्षर देखे**

Manufactured & Marketed by :

**INDOFIL INDUSTRIES LIMITED**

**Regd. Office :** Kalpataru Square, 4th Floor, Kondivita Road, Off Andheri-Kurla Road, Andheri (East), Mumbai - 400 059

04.23

# INDOFIL Z-78® ZINEB 75% WP

## (CONTACT FUNGICIDE)

It is wettable powder fungicides containing 75% w/w active ingredient Zineb and balance other relevant ingredients.

**Recommendation:**

| Crops | Common Name of Diseases | Dosage Per Hectare: A.I. Kg | Dosage Per Hectare: Formulation Kg | Dosage Per Hectare: Dilution in water (Ltr) | Waiting Period between last spray to harvest (in days) |
|---|---|---|---|---|---|
| Jowar | Red leaf spot | 1.125-1.5 | 1.5-2 | 750-1000 | |
| | Leaf Spot | 1.125-1.5 | 1.5-2 | 750-1000 | |
| | Leaf Blight | 1.125-1.5 | 1.5-2 | 750-1000 | |
| Paddy | Blast | 1.125-1.5 | 1.5-2 | 750-1000 | |
| Wheat | Rust | 1.125-1.5 | 1.5-2 | 750-1000 | |
| | Blight | 1.125-1.5 | 1.5-2 | 750-1000 | |
| Maize | Leaf Blight | 1.125-1.5 | 1.5-2 | 750-1000 | |
| Ragi (Bajra) | Blast | 1.125-1.5 | 1.5-2 | 750-1000 | |
| Tobacco | Leaf Spot | 1.125-1.5 | 1.5-2 | 750-1000 | |
| Turmeric | Leaf Spot | 1.125-1.5 | 1.5-2 | 750-1000 | |
| Onion | Downy Mildew | 1.125-1.5 | 1.5-2 | 750-1000 | |
| | Blight | 1.125-1.5 | 1.5-2 | 750-1000 | |
| Potato | Early Blight | 1.125-1.5 | 1.5-2 | 750-1000 | |
| | Late Blight | 1.125-1.5 | 1.5-2 | 750-1000 | |
| Tomato | Early Blight | 1.125-1.5 | 1.5-2 | 750-1000 | |
| | Late Blight | 1.125-1.5 | 1.5-2 | 750-1000 | |
| | Grey Leaf Mould | 1.125-1.5 | 1.5-2 | 750-1000 | |
| Chillies | Fruit Rot | 1.125-1.5 | 1.5-2 | 750-1000 | |
| | Leaf Spot | 1.125-1.5 | 1.5-2 | 750-1000 | |
| Brinjal | Blight | 1.125-1.5 | 1.5-2 | 750-1000 | |
| Cucurbits | Downy Mildew | 1.125-1.5 | 1.5-2 | 750-1000 | |
| | Anthracnose | 1.125-1.5 | 1.5-2 | 750-1000 | |
| | Leaf Spot | 1.125-1.5 | 1.5-2 | 750-1000 | |
| Cauliflower | Leaf Spot | 1.125-1.5 | 1.5-2 | 750-1000 | |
| Cumin | Early Blight | 1.125-1.5 | 1.5-2 | 750-1000 | |
| Apple | Scab | 1.125-1.5 | 1.5-2 | 750-1000 | |
| | Black Rot | 1.125-1.5 | 1.5-2 | 750-1000 | |
| Citrus | Greasy Spot | 1.125-1.5 | 1.5-2 | 750-1000 | |
| Cherries | Leaf Spot | 1.125-1.5 | 1.5-2 | 750-1000 | |
| Grapes | Downy Mildew | 1.125-1.5 | 1.5-2 | 750-1000 | |
| Guava | Fruit Rot | 1.125-1.5 | 1.5-2 | 750-1000 | |

**Direction of Use** : Mix the required quantity in a small quantity of water to thick uniform paste. Allowed to soak for 5-10 minutes. Add remaining quantity of water slowly while string continuously & make up the volume. Knapsack sprayer, foot sprayer, compression knapsack sprayer, compression knapsack battery sprayer. **Precaution :** Keep away from food stuffs and animals feed in cool dry place away from habitation. Do not smoke, chew, eat or drink during application. wash contaminated clothes and skin after application. Avoid skin contact. **Symptoms of Poisoning :** Nausea, vomitting, abdominal pain, diarrhoea, if swallowed. When inhaled as fine dust irritation to bronchail tubes and lungs tissues. **First Aid** : Induce vomitting by tickling the back of the throat, until vomit fluid is clear. In case of suspected poisioning call the physician. Wash contaminated skin with copious water. If eyes are contaminated, wash with plenty of clean water for minutes and refer to the doctor. **Antidote** : Treat symptomatically. **Disposal of Used Containers :** 1. The packages shall be broken and buried away from habitation. 2. The used packages shall not be left outside to prevent their reuse. 3. Packages as surplus material and washings should be disposed off in a safe manner so as to prevent environmental or water pollution. **Storage Conditions :** 1. The packages containing the fungicide shall be stored in separate rooms or premises away from the room or premises used for storing other articles or shall be kept in separate almirahs under lock and key depending upon the quantity and nature of the material. 2. The room or premises meant for storing the materials shall be well built, well lit, well ventilated and of sufficient dimension to prevent environmental pollution. **Chemical Composition** : Zineb Tech, Based on 80% a.i. 93.750% w/w, Wetting Agent/Suspending/Dispersing Agent Sodium salt of alkylaryl sulphonate, C.M.C & Glue. 2.000% w/w, Inert Carrier China Clay: 4.250% w/w, **Total** :100.000% w/w.

English