

మెటలోసోట్ ఇండోలైఫ్ వైటల్ ఎనర్జీ

| | |
|---------------------------|-----------|
| మొత్తం బోరాన్ (బొంబ) శాతం | 5.0 |
| బోరాన్ (బొంబ) కనీసం | 5.0 |
| పిఎస్ | 8.0-9.0 |
| నిట్రిజన్ (గ్రా/మి.లీ) | 1.10-1.23 |

అసంపూర్ణమైనదిగా ఉండటం వల్ల ఉపయోగించాలి. అసంపూర్ణమైనదిగా ఉండటం వల్ల ఉపయోగించాలి. అసంపూర్ణమైనదిగా ఉండటం వల్ల ఉపయోగించాలి.

మెటలోసోట్ ఇండోలైఫ్ వైటల్ ఎనర్జీ

| | |
|--|-----------|
| ఫాస్ఫోరస్ (బొంబ) శాతం | 5.0 |
| ఫాస్ఫోరస్ (బొంబ) కనీసం | 5.0 |
| నిట్రిజన్ గురుత్వకర్షణ శాతం (గ్రా/మి.లీ) | 1.10-1.23 |

అసంపూర్ణమైనదిగా ఉండటం వల్ల ఉపయోగించాలి. అసంపూర్ణమైనదిగా ఉండటం వల్ల ఉపయోగించాలి. అసంపూర్ణమైనదిగా ఉండటం వల్ల ఉపయోగించాలి.

మెటలోసోట్ ఇండోలైఫ్ వైటల్ ఎనర్జీ

| | |
|--|-----------|
| ఫాస్ఫోరస్ (బొంబ) శాతం | 5.0 |
| ఫాస్ఫోరస్ (బొంబ) కనీసం | 5.0 |
| నిట్రిజన్ గురుత్వకర్షణ శాతం (గ్రా/మి.లీ) | 1.10-1.23 |

అసంపూర్ణమైనదిగా ఉండటం వల్ల ఉపయోగించాలి. అసంపూర్ణమైనదిగా ఉండటం వల్ల ఉపయోగించాలి. అసంపూర్ణమైనదిగా ఉండటం వల్ల ఉపయోగించాలి.



FERTILIZER

Chelated Boron as Boron-Glycine (liquid)

Metalosate™

INDOLIFE™ VITAL ENERGY®

FERTILIZER Enriched with Boron
Albion™ & Metalosate™ are Trademarks of Albion Laboratories, Inc., U.S.A.
Balchem™ is a Trademark of Balchem Corporation, U.S.A.
Indolife Vital Energy® is a Trademark of INDOLFIL INDUSTRIES LIMITED, Mumbai, India.

ALBION™ BALCHEM™

PLANT NUTRITION

Fertilizer License No.: IQ/QC-5/Fert. Prmt/421-A/81920-941/2020
Manufactured by: ALBION LABORATORIES Inc., 67 South Main Street, Suite 200, Layton, Utah 84041, U.S.A. a wholly owned subsidiary of Balchem Corporation (U.S.A.).
Imported & Marketed by: INDOLFIL INDUSTRIES LIMITED

Regd. Office: Kalpataru Square, 4th Floor, Kondivita Road, Off Andheri-Kurja Road, Andheri (East), Mumbai-400 059

Re-Packed by: M/s Utsavi Agri Tech, Plot No. 3532, GIDC Industrial Estate, Panoli, Near Solvay Speciality Pvt. Ltd., Taluka: Ankleshwar, District: Bharuch, Gujarat - 394116

ఉత్పాదక: అల్బియన్ లాబొరేటరీస్, అయస్.ఎస్., 67 సౌత్ మెన్ స్ట్రీట్, స్యూ.ఎస్., లేటన్, యూటా 84041, యూ.ఎస్.ఎ., అల్బియన్ లాబొరేటరీస్ యొక్క పూర్తిగా నిర్వహించబడిన కుమార్తె సంస్థ.
అయిటల్ అండ్ మార్కెట్: ఇండోఫిల్ ఇండస్ట్రీస్ లిమిటెడ్
రెజి. ఆఫీస్: కల్పతారు స్కవైర్, 4వ ఫ్లోర్, కండవిటా రోడ్, ఆఫ్ ఆందెరి-కుర్జా రోడ్, ఆందెరి (ఈస్ట్), ముంబై-400 059
పున్: పిక్కెట్: M/s Utsavi Agri Tech, ప్లాట్ నెం. 3532, జి.డి.సి. ప్రొడ్యూక్స్ ఇండస్ట్రీయల్ ఎస్టేట్, పానోలి, గుజరాత్, భారత్. కే.బి.ఎస్., తాలూకా: ఆంకలేశ్వర్, జిల్లా: మరాఠ్వాడ, గుజరాత్-394116

For any complaint contact Customer Care Cell at Indofil Industries Limited, Kalpataru Square, 4th Floor, Kondivita Road, Off Andheri-Kurja Road, Andheri (East), Mumbai - 400 059.
Toll Free No.: 1800-120-003-004, email: customer.care@indofil.com



Imported & Marketed by:
INDOLFIL INDUSTRIES LIMITED
Regd. Office: Kalpataru Square, 4th Floor, Kondivita Road, Off Andheri-Kurja Road, Andheri (East), Mumbai - 400 059.
02.23

2) ద్వీకా పంపిణీ, 2వ పంపిణీ 20 రోజుల వ్యవస్థాపనలో అసంపూర్ణమైనదిగా ఉండటం వల్ల ఉపయోగించాలి. అసంపూర్ణమైనదిగా ఉండటం వల్ల ఉపయోగించాలి. అసంపూర్ణమైనదిగా ఉండటం వల్ల ఉపయోగించాలి.

2. ద్వీకా పంపిణీ, 2వ పంపిణీ 20 రోజుల వ్యవస్థాపనలో అసంపూర్ణమైనదిగా ఉండటం వల్ల ఉపయోగించాలి. అసంపూర్ణమైనదిగా ఉండటం వల్ల ఉపయోగించాలి. అసంపూర్ణమైనదిగా ఉండటం వల్ల ఉపయోగించాలి.

2. ద్వీకా పంపిణీ, 2వ పంపిణీ 20 రోజుల వ్యవస్థాపనలో అసంపూర్ణమైనదిగా ఉండటం వల్ల ఉపయోగించాలి. అసంపూర్ణమైనదిగా ఉండటం వల్ల ఉపయోగించాలి. అసంపూర్ణమైనదిగా ఉండటం వల్ల ఉపయోగించాలి.

Metalosate™ INDOLIFE Vital Energy®

Fertilizer Chelated Boron as Boron-Glycine (liquid)
Composition: (Percent by Weight)
Boron (as B) Minimum 5.0
pH 8.0 - 9.0
Specific Gravity (g/ml) 1.10-1.23
To be used where there is a recognized need. Do not exceed the appropriate application rate.
It is world's unique and an exclusive proprietary technology product specially designed and produced by Albion™. It improves strength of cell wall, pollen tube germination, increases flowering and retention and fruit development.
Recommendation: INDOLIFE Vital Energy® may be used on variety of Vegetables crops like Cabbage Cauliflower, Tomato, Cucurbits; Fruit crops like Grape, Citrus, Apple, Mango; Field crop like Rice, Maize, Wheat and many other field crops, cash crops, spices and condiments.
Time of Application: For Foliar Application
Crop Dose Time of spraying
Rice/ Maize 1-1.5ml/litre Flowering Stage
Wheat 1-1.5ml/litre Flowering stage
Vegetables 1-1.5ml/litre Flowering stage and other crops
Note: 1. We recommend minimum 200 litres of water / acre.
2. In long durations crops, need to go for > 2 sprays at an interval of 20 days depending on need.
Directions: For agriculture use only.
Shake well before use. Keep in a cool dry place, away from heat and open flame. Keep away from the reach of children.
Use of non-ionic surfactants may improve spray coverage.
Compatible with most of agrochemicals. Do not mix with Lime, Sulfur, Copper and Mineral oil. It's advisable to test compatibility before mixing. Do not apply undiluted.
First aid measures: There are not specific hazards known to be associated with this product, although precautions should be taken to avoid unnecessary contact with skin, eyes and mouth. Eyes: Irrigate thoroughly with water at least 10 minutes. If any discomfort persists obtain medical attention. Skin: Wash skin thoroughly with water. Remove contaminated clothing and wash before reuse. In severe cases obtain medical attention. Ingestion: Wash out mouth thoroughly with water. Obtain medical attention. Inhalation: Remove from exposure, rest and keep warm. In severe cases obtain medical attention.
Disclaimer: As use of this product is beyond our control, we do not take any responsibility other than uniform quality of the product.

মেটালোসেট ইন্ডোলাইফ বায়টল এনর্জি

Fertilizer Chelated Boron as Boron-Glycine (liquid)
Composition: (Percent by Weight)
Boron (as B) Minimum 5.0
pH 8.0 - 9.0
Specific Gravity (g/ml) 1.10-1.23
To be used where there is a recognized need. Do not exceed the appropriate application rate.
It is world's unique and an exclusive proprietary technology product specially designed and produced by Albion™. It improves strength of cell wall, pollen tube germination, increases flowering and retention and fruit development.
Recommendation: INDOLIFE Vital Energy® may be used on variety of Vegetables crops like Cabbage Cauliflower, Tomato, Cucurbits; Fruit crops like Grape, Citrus, Apple, Mango; Field crop like Rice, Maize, Wheat and many other field crops, cash crops, spices and condiments.
Time of Application: For Foliar Application
Crop Dose Time of spraying
Rice/ Maize 1-1.5ml/litre Flowering Stage
Wheat 1-1.5ml/litre Flowering stage
Vegetables 1-1.5ml/litre Flowering stage and other crops
Note: 1. We recommend minimum 200 litres of water / acre.
2. In long durations crops, need to go for > 2 sprays at an interval of 20 days depending on need.
Directions: For agriculture use only.
Shake well before use. Keep in a cool dry place, away from heat and open flame. Keep away from the reach of children.
Use of non-ionic surfactants may improve spray coverage.
Compatible with most of agrochemicals. Do not mix with Lime, Sulfur, Copper and Mineral oil. It's advisable to test compatibility before mixing. Do not apply undiluted.
First aid measures: There are not specific hazards known to be associated with this product, although precautions should be taken to avoid unnecessary contact with skin, eyes and mouth. Eyes: Irrigate thoroughly with water at least 10 minutes. If any discomfort persists obtain medical attention. Skin: Wash skin thoroughly with water. Remove contaminated clothing and wash before reuse. In severe cases obtain medical attention. Ingestion: Wash out mouth thoroughly with water. Obtain medical attention. Inhalation: Remove from exposure, rest and keep warm. In severe cases obtain medical attention.
Disclaimer: As use of this product is beyond our control, we do not take any responsibility other than uniform quality of the product.

মেটালোসেট ইন্ডোলাইফ বায়টল এনর্জি

Fertilizer Chelated Boron as Boron-Glycine (liquid)
Composition: (Percent by Weight)
Boron (as B) Minimum 5.0
pH 8.0 - 9.0
Specific Gravity (g/ml) 1.10-1.23
To be used where there is a recognized need. Do not exceed the appropriate application rate.
It is world's unique and an exclusive proprietary technology product specially designed and produced by Albion™. It improves strength of cell wall, pollen tube germination, increases flowering and retention and fruit development.
Recommendation: INDOLIFE Vital Energy® may be used on variety of Vegetables crops like Cabbage Cauliflower, Tomato, Cucurbits; Fruit crops like Grape, Citrus, Apple, Mango; Field crop like Rice, Maize, Wheat and many other field crops, cash crops, spices and condiments.
Time of Application: For Foliar Application
Crop Dose Time of spraying
Rice/ Maize 1-1.5ml/litre Flowering Stage
Wheat 1-1.5ml/litre Flowering stage
Vegetables 1-1.5ml/litre Flowering stage and other crops
Note: 1. We recommend minimum 200 litres of water / acre.
2. In long durations crops, need to go for > 2 sprays at an interval of 20 days depending on need.
Directions: For agriculture use only.
Shake well before use. Keep in a cool dry place, away from heat and open flame. Keep away from the reach of children.
Use of non-ionic surfactants may improve spray coverage.
Compatible with most of agrochemicals. Do not mix with Lime, Sulfur, Copper and Mineral oil. It's advisable to test compatibility before mixing. Do not apply undiluted.
First aid measures: There are not specific hazards known to be associated with this product, although precautions should be taken to avoid unnecessary contact with skin, eyes and mouth. Eyes: Irrigate thoroughly with water at least 10 minutes. If any discomfort persists obtain medical attention. Skin: Wash skin thoroughly with water. Remove contaminated clothing and wash before reuse. In severe cases obtain medical attention. Ingestion: Wash out mouth thoroughly with water. Obtain medical attention. Inhalation: Remove from exposure, rest and keep warm. In severe cases obtain medical attention.
Disclaimer: As use of this product is beyond our control, we do not take any responsibility other than uniform quality of the product.

মেটালোসেট ইন্ডোলাইফ বায়টল এনর্জি

Fertilizer Chelated Boron as Boron-Glycine (liquid)
Composition: (Percent by Weight)
Boron (as B) Minimum 5.0
pH 8.0 - 9.0
Specific Gravity (g/ml) 1.10-1.23
To be used where there is a recognized need. Do not exceed the appropriate application rate.
It is world's unique and an exclusive proprietary technology product specially designed and produced by Albion™. It improves strength of cell wall, pollen tube germination, increases flowering and retention and fruit development.
Recommendation: INDOLIFE Vital Energy® may be used on variety of Vegetables crops like Cabbage Cauliflower, Tomato, Cucurbits; Fruit crops like Grape, Citrus, Apple, Mango; Field crop like Rice, Maize, Wheat and many other field crops, cash crops, spices and condiments.
Time of Application: For Foliar Application
Crop Dose Time of spraying
Rice/ Maize 1-1.5ml/litre Flowering Stage
Wheat 1-1.5ml/litre Flowering stage
Vegetables 1-1.5ml/litre Flowering stage and other crops
Note: 1. We recommend minimum 200 litres of water / acre.
2. In long durations crops, need to go for > 2 sprays at an interval of 20 days depending on need.
Directions: For agriculture use only.
Shake well before use. Keep in a cool dry place, away from heat and open flame. Keep away from the reach of children.
Use of non-ionic surfactants may improve spray coverage.
Compatible with most of agrochemicals. Do not mix with Lime, Sulfur, Copper and Mineral oil. It's advisable to test compatibility before mixing. Do not apply undiluted.
First aid measures: There are not specific hazards known to be associated with this product, although precautions should be taken to avoid unnecessary contact with skin, eyes and mouth. Eyes: Irrigate thoroughly with water at least 10 minutes. If any discomfort persists obtain medical attention. Skin: Wash skin thoroughly with water. Remove contaminated clothing and wash before reuse. In severe cases obtain medical attention. Ingestion: Wash out mouth thoroughly with water. Obtain medical attention. Inhalation: Remove from exposure, rest and keep warm. In severe cases obtain medical attention.
Disclaimer: As use of this product is beyond our control, we do not take any responsibility other than uniform quality of the product.

মেটালোসেট ইন্ডোলাইফ বায়টল এনর্জি

Fertilizer Chelated Boron as Boron-Glycine (liquid)
Composition: (Percent by Weight)
Boron (as B) Minimum 5.0
pH 8.0 - 9.0
Specific Gravity (g/ml) 1.10-1.23
To be used where there is a recognized need. Do not exceed the appropriate application rate.
It is world's unique and an exclusive proprietary technology product specially designed and produced by Albion™. It improves strength of cell wall, pollen tube germination, increases flowering and retention and fruit development.
Recommendation: INDOLIFE Vital Energy® may be used on variety of Vegetables crops like Cabbage Cauliflower, Tomato, Cucurbits; Fruit crops like Grape, Citrus, Apple, Mango; Field crop like Rice, Maize, Wheat and many other field crops, cash crops, spices and condiments.
Time of Application: For Foliar Application
Crop Dose Time of spraying
Rice/ Maize 1-1.5ml/litre Flowering Stage
Wheat 1-1.5ml/litre Flowering stage
Vegetables 1-1.5ml/litre Flowering stage and other crops
Note: 1. We recommend minimum 200 litres of water / acre.
2. In long durations crops, need to go for > 2 sprays at an interval of 20 days depending on need.
Directions: For agriculture use only.
Shake well before use. Keep in a cool dry place, away from heat and open flame. Keep away from the reach of children.
Use of non-ionic surfactants may improve spray coverage.
Compatible with most of agrochemicals. Do not mix with Lime, Sulfur, Copper and Mineral oil. It's advisable to test compatibility before mixing. Do not apply undiluted.
First aid measures: There are not specific hazards known to be associated with this product, although precautions should be taken to avoid unnecessary contact with skin, eyes and mouth. Eyes: Irrigate thoroughly with water at least 10 minutes. If any discomfort persists obtain medical attention. Skin: Wash skin thoroughly with water. Remove contaminated clothing and wash before reuse. In severe cases obtain medical attention. Ingestion: Wash out mouth thoroughly with water. Obtain medical attention. Inhalation: Remove from exposure, rest and keep warm. In severe cases obtain medical attention.
Disclaimer: As use of this product is beyond our control, we do not take any responsibility other than uniform quality of the product.